

IDENTITIES BETWEEN MAHLER MEASURES

FERNANDO RODRIGUEZ-VILLEGAS

Department of Mathematics
University of Texas at Austin
Austin, TX 78712

September 2000

The purpose of this short note is to give a proof of the following identity between (logarithmic) Mahler measures

$$(1) \quad m(y^2 + 2xy + y - x^3 - 2x^2 - x) = \frac{5}{7} m(y^2 + 4xy + y - x^3 + x^2),$$

which is one of many examples that arise from the comparison of Mahler measures and special values of L -functions [Bo], [De], [RV]. Let us recall that the *logarithmic Mahler measure* of a Laurent polynomial $P \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ is defined as

$$(2) \quad m(P) = \int_0^1 \cdots \int_0^1 \log |P(e^{2\pi i \theta_1}, \dots, e^{2\pi i \theta_n})| d\theta_1 \cdots d\theta_n.$$

The conjecture of Bloch–Beilinson [Be], [BG] for elliptic curves predicts that both sides of (1) are rationally related to $L'(E, 0)$ (and hence to each other), where E is the elliptic curve of conductor 37

$$(3) \quad E : \quad y^2 + y = x^3 - x,$$

Support for this work was provided in part by a grant of the NSF and by a Sloan Research Fellowship.

and $L(E, s)$ is its L -function. More precisely, we expect that the two numbers a and b defined by

$$\begin{aligned} m(y^2 + 2xy + y - x^3 - 2x^2 - x) &= a L'(E, 0), \\ m(y^2 + 4xy + y - x^3 + x^2) &= b L'(E, 0) \end{aligned}$$

are rational. A proof of this fact is not without reach but will not be attempted here, we will prove instead that $a/b = 5/7$.

1. Computing in $K_2(E)$

We first recall the definition of the group $K_2(A)$ of an elliptic curve A . Given a field F the group $K_2(F)$ can be defined as $F^* \otimes F^*$ modulo the Steinberg relations $x \otimes (1 - x)$ for $x \neq 0, 1$ in F .

Given a discrete valuation v on F with maximal ideal $\mathcal{M}$ and residue field k we have the *tame symbol* at v defined by

$$(x, y)_v \equiv (-1)^{v(x)v(y)} \frac{x^{v(y)}}{y^{v(x)}} \pmod{\mathcal{M}},$$

which determines a homomorphism

$$\lambda_v : K_2(F) \longrightarrow k^*$$

For an elliptic curve A defined over $\mathbb{Q}$ we let $K_2(A)$ be the elements of $K_2(\mathbb{Q}(A))$ annihilated by all λ_v with v the valuations associated to $\overline{\mathbb{Q}}$ points of A .

Our E appears as a fiber in several of Boyd's families of elliptic curves (see [Bo], [RV] for a discussion of these families). For example, in its original form $y^2 + y = x^3 - x$, but also as the two Weierstrass equations

$$(4) \quad E_1 : \quad y_1^2 + 4x_1y_1 + y_1 = x_1^3 - x_1^2$$

and

$$(5) \quad E_2 : \quad y_2^2 + 2x_2y_2 + y_2 = x_2^3 + 2x_2^2 + x_2.$$

It is easy to check that

$$(6) \quad \begin{aligned} x_1 &= x - 1 \\ y_1 &= y - 2x + 2 \end{aligned}$$

and

$$(7) \quad \begin{aligned} x_2 &= x - 1 \\ y_2 &= -x + y + 1 \end{aligned}$$

give isomorphisms

$$E \simeq E_1, \quad E \simeq E_2.$$

It follows from [RV] therefore, that some integer multiple of each of

$$\xi = \{x, y\}, \quad \xi_1 = \{x_1, y_1\}, \quad \xi_2 = \{x_2, y_2\}$$

is in $K_2(E)$.

The divisors of the six functions x, y, x_1, y_1, x_2, y_2 are supported on $E(\mathbb{Q})$, which is generated by the point P with $x = 0, y = 0$. More precisely, we have

$$(8) \quad \begin{aligned} (x) &= [P] + [-P] - 2[O] \\ (y) &= [P] + [2P] + [-3P] - 3[O] \\ (x_1) &= [2P] + [-2P] - 2[O] \\ (y_1) &= 2[2P] + [-4P] - 3[O] \\ (x_2) &= [-2P] + [2P] - 2[O] \\ (y_2) &= [2P] + 2[-P] - 3[O] \end{aligned}$$

where $[O]$ denotes the point at infinity on E .

Given a pair of functions f and g on E with divisors supported on $E(\mathbb{Q})$

$$(f) = \sum_{n \in \mathbb{Z}} a_n [nP], \quad (g) = \sum_{n \in \mathbb{Z}} b_n [nP]$$

we define

$$(9) \quad (f) \diamond (g) = \sum_{m,n} a_n b_m [(n-m)P],$$

which we will view as an element of

$$\mathbb{Z}[E(\mathbb{Q})]^- = \mathbb{Z}[E(\mathbb{Q})]/\sim,$$

where $\sim$ is the equivalence relation determined by

$$[-nP] \sim -[nP], \quad n \in \mathbb{Z}.$$

We may and will represent elements of $\mathbb{Z}[E(\mathbb{Q})]^-$ as vectors $[a_1, a_2, \dots]$ with $a_i \in \mathbb{Z}$ almost all zero where

$$[a_1, a_2, \dots] \longleftrightarrow \sum_{n=1}^{\infty} a_n [nP]$$

In fact, we will only consider elements where $a_n = 0$ for $n > 6$ and hence simply write $[a_1, \dots, a_6]$.

We now compute

$$(10) \quad \begin{aligned} (x) \diamond (y) &= [1, 2, -3, 1, 0, 0] \\ (x_1) \diamond (y_1) &= [0, 5, 0, -4, 0, 1] \\ (x_2) \diamond (y_2) &= [-6, 2, 2, -1, 0, 0]. \end{aligned}$$

On the other hand, we also find

$$(11) \quad \begin{aligned} (-y) \diamond (1+y) &= [-8, -7, 8, 1, 0, -1] \\ (x-y) \diamond (1-x+y) &= [-9, 5, -5, 5, 0, -1] \end{aligned}$$

and verify easily that

$$(12) \quad \begin{aligned} 7(x) \diamond (y) + (x_1) \diamond (y_1) &= -2(-y) \diamond (1+y) + (x-y) \diamond (1-x+y) \\ 5(x) \diamond (y) + (x_2) \diamond (y_2) &= -(-y) \diamond (1+y) + (x-y) \diamond (1-x+y). \end{aligned}$$

2. The regulator

Let

$$(13) \quad r : K_2(E) \longrightarrow \mathbb{R}$$

be the regulator map. It can be defined as follows. If f, g are two non-constant functions on E with $\{f, g\} \in K_2(E)$ then

$$(14) \quad r(\{f, g\}) = \int_{\gamma} \eta(f, g),$$

where

$$(15) \quad \eta(f, g) = \log |f| \, d \arg g - \log |g| \, d \arg f$$

and γ is a closed path not going through poles or zeroes of f or g which generates the subgroup $H_1(E, \mathbb{Z})^-$ of $H_1(E, \mathbb{Z})$ where complex conjugation acts by -1 , properly oriented. The fact that the integral only depends on the homology class of γ is a consequence of $\{f, g\} \in K_2(E)$, see [RV] for details. (However, note that in [RV] we inaccurately said γ should generate the cycles fixed by complex conjugation; we take the opportunity to correct this.)

The regulator may also be expressed in terms of the elliptic dilogarithm [BG], [Za]

$$\mathcal{L} : E(\mathbb{C}) \longrightarrow \mathbb{R}.$$

In our context, this works as follows. We extend it by linearity to $\mathbb{Z}[E(\mathbb{Q})]$ and since $\mathcal{L}$ is odd it actually gives a map

$$(16) \quad \mathcal{L} : \mathbb{Z}[E(\mathbb{Q})]^- \longrightarrow \mathbb{R}.$$

If f, g are two non-constant functions on E with divisors supported on $E(\mathbb{Q})$ and such that $\{f, g\} \in K_2(E)$ then

$$(17) \quad r(\{f, g\}) = c \, \mathcal{L}((f) \diamond (g)),$$

for some explicit non-zero constant c , which is not relevant for our purposes. In particular, in the case that $g = 1 - f$

$$(18) \quad \mathcal{L}((f) \diamond (1 - f)) = 0.$$

The above discussion extends naturally to $K_2(E) \otimes \mathbb{Q}$, which contains ξ, ξ_1 and ξ_2 .

It follows from (12) therefore, that

$$(19) \quad \begin{aligned} r(\xi_1) &= -7r(\xi) \\ r(\xi_2) &= -5r(\xi) . \end{aligned}$$

3. The regulator and Mahler's measure

In [RV] we showed that if $P_k(x, y) = 0$ is one of Boyd's families of elliptic curves and k is such that P_k does not vanish on the torus $|x| = |y| = 1$ then

$$(20) \quad r(\{x, y\}) = c_k \pi \, m(P_k)$$

for some nonzero integer c_k . We will now make this precise for

$$P_k(x, y) = y^2 - kxy + y - x^3 + x^2 .$$

We consider the region $\mathcal{K}$ of $k \in \mathbb{C}$ such that P_k vanishes somewhere on the torus. It is the image of the torus under the rational map

$$(21) \quad R : \quad (x, y) \mapsto \frac{y^2 + y - x^3 + x^2}{xy} .$$

We can get a pretty good idea of what $\mathcal{K}$ looks like by graphing the image of a grid under $(\theta_1, \theta_2) \mapsto R(e^{2\pi i \theta_1}, e^{2\pi i \theta_2})$. Dividing the square $0 \leq \theta_1 < 1, 0 \leq \theta_2 < 1$ in 40 equal parts we obtain

FIGURE 1. Region $\mathcal{K}$

It is not hard to verify directly that the boundary of $\mathcal{K}$ meets the real axis at $k = -4$ and $k = 2$.

If $k \notin \mathcal{K}$ then as x moves counterclockwise on the circle $|x| = 1$ one root $y_1(x)$ of $P_k(x, y) = 0$ satisfies $|y_1(x)| < 1$ and the other $y_2(x)$ satisfies $|y_2(x)| > 1$ and in particular $y_1(x)$ and $y_2(x)$ do not meet. To see this, note

that when $x = 1$ the roots are 0 and $k - 1$. Hence, for $|k|$ large these roots are one inside and the other outside the unit circle. The claim follows since the roots depends continuously on k . We let σ_k be the resulting smooth closed path $(x, y_1(x))$ on the elliptic curve E_k determined by $P_k(x, y) = 0$.

Using Jensen's formula we find that

$$m(P_k) = \frac{1}{2\pi i} \int_{\sigma_k} \log |y| \frac{dx}{x}$$

and note that since $|x| = 1$ on σ_k we can write this identity as

$$(22) \quad m(P_k) = \frac{1}{2\pi} \int_{\sigma_k} \eta(x, y).$$

We now show that for real and $k \notin \mathcal{K}$ the homology class of σ_k generates $H_1(E_k, \mathbb{Z})^-$. We complete the square and write $P_k = (2y - kx + 1)^2 - f(x)$, where $f(x) = 4x^3 + (k^2 - 4)x^2 - 2kx + 1$. The discriminant $\Delta(k) = k^4 - k^3 - 8k^2 + 36k - 1$ of f has two real roots $\alpha = -3.7996\dots$ and $\beta = .3305\dots$. Hence, for $k < \alpha$ or $k > \beta$, $\Delta(k) > 0$ and f has three real roots $e_1 < e_2 < e_3$. As $|k|$ increases the roots of f tend to $e_1 = -\infty$ and $e_2 = e_3 = 0$ and by continuity the circle $|x| = 1$ encircles e_2 and e_3 once. Since f is negative in the interval $e_2 < x < e_3$ the period

$$\int_{\sigma_k} \frac{dx}{2y - kx + 1}$$

is purely imaginary and our claim follows.

Combined with (14) and (22) this proves that in fact

$$(23) \quad r(\{x, y\}) = \pm 2\pi m(P_k), \quad k \in \mathbb{R}, \quad k \notin \mathcal{K}.$$

By continuity (23) also holds for $k = -4$ and $k = 2$, which are on the boundary of $\mathcal{K}$. In particular, in the notation of §2, we obtain the identity

$$(24) \quad r(\xi_1) = \pm 2\pi m(y^2 + 4xy + y - x^3 + x^2).$$

A completely analogous analysis yields

$$(25) \quad r(\xi_2) = \pm 2\pi m(y^2 + 2xy + y - x^3 - 2x^2 - x)$$

(and again $k = -2$ is on the boundary of the corresponding set $\mathcal{K}$). Putting together (19), (24) and (25) (and a simple check for the right sign) we obtain (1).

Remarks 1. We should point out that we do not expect $m(y^2 + y - x^3 + x)$ to be rationally related to either side of (1) (and numerically it indeed does not appear to be). The reason is that $y^2 + y - x^3 + x$ vanishes on the torus and in fact $k = 0$ is in the interior of the region $\mathcal{K}$ corresponding to the Boyd family $y^2 - kxy + y - x^3 + x$. Hence the analogue of (22) gives the integral of $\eta(x, y)$ on a non-closed cycle.

2. One can prove in a similar way an identity relating either side of (1) with $m(y^2 + 2xy + y - x^3 + x^2)$.

REFERENCES

- [Be] A. Beilinson, *Higher regulators of modular curves*, Applications of algebraic K -theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983), Contemp. Math., vol. **55**, Amer. Math. Soc., Providence, R.I., 1986, pp. 1–34.
- [Bo] D. W. Boyd, *Mahler’s measure and special values of L -functions*, Experiment. Math. **7** (1998), 37–82.
- [BG] S. Bloch & D. Grayson, *K_2 and L -functions of elliptic curves: Computer Calculations*, Contemp. Math. **55** (1986), 79–88.
- [De] C. Deninger, *Deligne periods of mixed motives, K -theory and the entropy of certain $\mathbb{Z}^n$ -actions*, J. Amer. Math. Soc. **10** (1997), 259–281.
- [RV] F. Rodriguez Villegas, *Modular Mahler measures I*, Topics in Number Theory (S.D. Ahlgren, G.E. Andrews & K. Ono, ed.), Kluwer, Dordrecht, 1999, pp. 17–48.
- [Z] D. Zagier, *The Bloch–Wigner–Ramakrishnan polylogarithm function*, Math. Ann. **286** (1990), 613–624.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN

PREPRINT

E-mail address: villegas@math.utexas.edu